

環境と子供の健康に関するコホート研究（北海道スタディからの論文）
(PFOS, PFOA等)

1. Inoue K, Okada F, Ito R, Kato S, Sasaki S, Nakajima S, Uno A, Saijo Y, Sata F, Yoshimura Y, Kishi R, Nakazawa H: Perfluorooctane sulfonate (PFOS) and related perfluorinated compounds in human maternal and cord blood samples: assessment of PFOS exposure in a susceptible population during pregnancy. **Environ Health Perspect.** **112(11):1204-1207, 2004**
2. Washino N, Saijo Y, Sasaki S, Kato S, Ban S, Konishi K, Ito R, Nakata A, Iwasaki Y, Saito K, Nakazawa H, Kishi R: Correlations between prenatal exposure to perfluorinated chemicals and reduced fetal growth. **Environ Health Perspect.** **117(4):660-667, 2009**
3. Okada E, Sasaki S, Saijo Y, Washino N, Miyashita C, Kobayashi S, Konishi K, Ito YM, Ito R, Nakata A, Iwasaki Y, Saito K, Nakazawa H, Kishi R: Prenatal exposure to perfluorinated chemicals and relationship with allergies and infectious diseases in infants. **Environ Res.** **112:118-125, 2012**
4. Okada E., Kashino I., Matsuura H., Sasaki S., Miyashita C., Yamamoto J., Ikeno T., Ito M. Y., Matsumura T., Tamakoshi A., Kishi R.; Temporal trends of perfluoroalkyl acids in plasma samples of pregnant women in Hokkaido, Japan, 2003-2011. **Environ Int.** **60 89-96, 2013.**
5. Okada E., Sasaki S., Kashino I., Matsuura H., Miyashita C., Kobayashi S., Itoh K., Ikeno T., Tamakoshi A., Kishi R.; Prenatal exposure to perfluoroalkyl acids and allergic diseases in early childhood. **Environ Int.** **65 127-134, 2014.**
6. Kishi R., Nakajima T., Goudarzi H., Kobayashi S., Sasaki S., Okada E., Miyashita C., Ito S., Araki A., Ikeno T., Iwasaki Y. and Nakazawa H.; The Association of Prenatal Exposure to Perfluorinated Chemicals with Maternal Essential and Long-Chain Polyunsaturated Fatty Acids during Pregnancy and the Birth Weight of their Offspring: The Hokkaido Study, **Environ Health Perspect.** **123, (10):1038-45, 2015.**

7. Goudarzi H., Miyashita C., Okada E., Kashino I., Kobayashi S., Chen C. J., Ito S., Araki A., Matsuura H., Ito Y. M., Kishi R.; Effects of prenatal exposure to perfluoroalkyl acids on prevalence of allergic diseases among 4-year-old children. **Environ Int.** **94** 124-132, 2016.
8. Goudarzi H., Nakajima S., Ikeno T., Sasaki S., Kobayashi S., Miyashita C., Ito S., Araki A., Nakazawa H., Kishi R.; Prenatal exposure to perfluorinated chemicals and neurodevelopment in early infancy: The Hokkaido Study. **Sci Total Environ.** **541** 1002-1010, 2016.
9. Itoh S., Araki A., Mitsui T., Miyashita C., Goudarzi H., Sasaki S., Cho K., Nakazawa H., Iwasaki Y., Shinohara N., Nonomura K., Kishi R.; Association of perfluoroalkyl substances exposure in utero with reproductive hormone levels in cord blood in the Hokkaido Study on Environment and Children's Health. **Environ Int.** **94** 51-59, 2016.
10. Kato S., Itoh S., Yuasa M., Baba T., Miyashita C., Sasaki S., Nakajima S., Uno A., Nakazawa H., Iwasaki Y., Okada E., Kishi R.; Association of perfluorinated chemical exposure in utero with maternal and infant thyroid hormone levels in the Sapporo cohort of Hokkaido Study on the Environment and Children's Health. **Environ Health Prev Med.** **21**, 334-344, 2016.
11. Minatoya M., Itoh S., Miyashita C., Araki A., Sasaki S., Miura R., Goudarzi H., Iwasaki Y., Kishi R.; Association of prenatal exposure to perfluoroalkyl substances with cord blood adipokines and birth size: The Hokkaido Study on environment and children's health. **Environ Res.** **156** 175-182, 2017.
12. Kobayashi S., Azumi K., Goudarzi H., Araki A., Miyashita C., Kobayashi S., Itoh S., Sasaki S., Ishizuka M., Nakazawa H., Ikeno T., Kishi R.; Effects of prenatal perfluoroalkyl acid exposure on cord blood IGF2/H19 methylation and ponderal index: the Hokkaido study. **J Expo Sci Environ Epidemiol.** **27** (3):251-259, 2017.
13. Goudarzi H., Miyashita C., Okada E., Kashino I., Chen C. J., Ito S., Araki A., Kobayashi S., Matsuura H., Kishi R.; Prenatal exposure to perfluoroalkyl acids and prevalence of infectious diseases up to 4years of age. **Environ Int.** **104** 132-138, 2017.

14. Goudarzi H., Araki A., Itoh S., Sasaki S., Miyashita C., Mitsui T., Nakazawa H., Nonomura K., Kishi R.; The Association of Prenatal Exposure to Perfluorinated Chemicals with Glucocorticoid and Androgenic Hormones in Cord Blood Samples: The Hokkaido Study. **Environ Health Perspect.** **125 (1):111-118, 2017.**
15. Miura R., Araki A., Miyashita C., Kobayashi Su., Kobayashi Sa., Wang SL., Chen CH., Miyake K., Ishizuka M., Iwasaki Y., YM. Ito, Kubota T., Kishi R.; An epigenome-wide study of cord blood DNA methylations in relation to prenatal perfluoroalkyl substance exposure: the Hokkaido study. **Environ Int.** **115 21-28, 2018.**
16. Tsai M.-S., Miyashita C., Araki A., Itoh S., Ait Bamai Y., Goudarzi H., Okada E., Kashino I., Matsuura H., Kishi R.; Determinants and temporal trends of perfluoroalkyl substances in pregnant women: The Hokkaido Study on Environment and Children's Health. **Int. J. Environ. Res. Public Health** **15 (5) 989, 2018.**
17. Itoh S., Araki A., Miyashita C., Yamazaki K., Goudarzi H., Minatoya M., Ait Bamai Y., Kobayashi S., Okada E., Kashino I., Yuasa M., Baba T., Kishi R., Association between perfluoroalkyl substance exposure and thyroid hormone/thyroid antibody levels in maternal and cord blood: The Hokkaido Study., **Environ Int,** **133, Part A, 105139, 2019.**
18. Kashino I., Sasaki S., Okada E., Matsuura H., Goudarzi H., Miyashita C., Okada E., Ito Y. M., Araki A., Kishi R.; Prenatal exposure to 11 perfluoroalkyl substances and fetal growth: A large-scale, prospective birth cohort study. **Environ Int.** **136 105355, 2020.**
19. Ait Bamai Y., Goudarzi H., Araki A., Okada E., Kashino I., Miyashita C., Kishi R.; Effect of prenatal exposure to per- and polyfluoroalkyl substances on childhood allergies and common infectious diseases in children up to age 7 years: The Hokkaido study on environment and children's health. **Environ Int.** **143 105979, 2020.**
20. Kobayashi S., Sata F., Ikeda-Araki A., Miyashita C., Itoh S., Goudarzi H., Iwasaki Y., Mitsui T., Moriya K., Shinohara N., Cho K., Kishi R., Associations among maternal perfluoroalkyl substance levels, fetal sex-hormone enzymatic gene polymorphisms, and fetal sex hormone levels in the Hokkaido study. **Reprod. Toxicol.** **105:221-231, 2021 Oct.**

21. Kobayashi S., Sata F., Goudarzi H., Araki A., Miyashita C., Sasaki S., Okada E., Iwasaki Y., Nakajima T., Kishi R., Associations among perfluorooctanesulfonic/perfluorooctanoic acid levels, nuclear receptor gene polymorphisms, and lipid levels in pregnant women in the Hokkaido study. **Sci. Rep.** **11;11(1), 9994, 2021 May.**
22. Itoh S., Yamazaki K., Suyama S., Ikeda-Araki A., Miyashita C., Ait Bamai Y., Kobayashi S., Masuda H., Yamaguchi T., Goudarzi H., Okada E., Kashino I., Saito T., Kishi R., The association between prenatal perfluoroalkyl substance exposure and symptoms of attention-deficit/hyperactivity disorder in 8-year-old children and the mediating role of thyroid hormones in the Hokkaido study. **Environ Int.** **159: 107026. 2022 Jan.**
23. Kobayashi S., Sata F., Ikeda-Araki A., Miyashita C., Goudarzi H., Iwasaki Y., Nakajima T., Kishi R., Relationships between maternal perfluoroalkyl substance levels, polymorphisms of receptor genes, and adverse birth outcomes in the Hokkaido birth cohort study, Japan. **Reprod. Toxicol.**, **107: 112-122, 2022 Jan.**
24. Nishimura Y., Moriya K., Kobayashi S., Ikeda-Araki A., Sata F., Mitsui T., Itoh S., Miyashita C., Cho K., Kon M., Nakamura M., Kitta T., Murai S., Kishi R., Shinohara N.. Association of exposure to prenatal perfluoroalkyl substances and estrogen receptor 1 polymorphisms with the second to fourth digit ratio in school-aged children: the Hokkaido Study. **Reprod. Toxicol.** **109, 10-18, 2022 Apr.**